

# A homological model for the coloured Jones polynomials

CRISTINA PALMER-ANGHEL

## Abstract

In 1991, Reshetikhin and Turaev defined a method that starts with a quantum group and leads to link invariants. This construction is purely algebraic and combinatorial. The coloured Jones polynomials  $J_N(L, q)$  are a family of quantum invariants for links constructed using this method from the representation theory of the quantum group  $U_q(sl(2))$ . The first invariant from this sequence, corresponding to the parameter  $N = 2$  is the original Jones polynomial  $J(q)$ . This is a quantum invariant, but it can be characterized also by skein relations. We will describe a homological model for all coloured Jones polynomials.

R. Lawrence defined a sequence of representations of the braid groups, using the homology of a certain covering of a configuration space. From these homological representations, Bigelow [2] and Lawrence [5] described the original Jones polynomial as a graded intersection pairing in a covering of a configuration space of the punctured disk, using the skein nature of the invariant for the proof.

In 2012, Kohno ([4], [3]) proved a deep connection between the representations of the braid group on the highest weight spaces of  $U_q(sl(2))$ -modules and the Lawrence representations. We present a homological model for  $J_N(L, q)$ , using this identification and the quantum nature of the invariant for the proof. We show that any coloured Jones polynomial can be described as a graded intersection pairing between two homology classes on a covering of the configuration space of the punctured disc.

This work is available at [1].

## References

- [1] Cristina A-M Anghel - A homological model for the coloured Jones polynomials. <https://arxiv.org/abs/1712.04873> , 2017.
- [2] Stephen Bigelow - A homological definition of the Jones polynomial. In Invariants of knots and 3-manifolds (Kyoto, 2001), volume 4 of Geom. Topol. Monogr., pages 29-41. Geom. Topol. Publ., Coventry, 2002.
- [3] Tetsuya Ito - Reading the dual Garside length of braids from homological and quantum representations. Comm. Math. Phys., 335(1):345-367, 2015.
- [4] Toshitake Kohno- Quantum and homological representations of braid groups. Configuration Spaces - Geometry, Combinatorics and Topology, Edizioni della Normale (2012), 355-372.
- [5] R. J. Lawrence - A functorial approach to the one-variable Jones polynomial. J. Differential Geom., 37(3):689-710, 1993.

(Cristina Palmer-Anghel) PARIS DIDEROT UNIVERSITY  
e-mail: [cristina.anghel@imj-prg.fr](mailto:cristina.anghel@imj-prg.fr)